



The Oklahoma Poison Control Center

940 N.E. 13th Street, Oklahoma City, Oklahoma 73104

Office Phone: (405) 271-5062 Fax: (405) 271-1816

www.oklahomapoison.org



Antidote Survey

April 16, 2007

If these antidotes are located in areas other than the main pharmacy (i.e. ER, satellites, etc.), please duplicate this survey to be filled out in the additional locations. Please indicate the location of the antidotes by checking one of the following boxes.

main pharmacy ER satellite pharmacy other _____

Antidote	Concentration and Size	Number of Units/Vials
Antivenin, polyvalent FAB (CroFab®; IV)	Single use vial	
Antivenin, Latrodectus mactans (black widow; IV)	6000 units/vial	
Calcium chloride	10%. In 10 mL amps	
	10%. In 10mL vials	
	10%. In 10mL syringes	
Calcium disodium EDTA (Versenate®)	200 mg/mL. In 5 mL amps.	
Calcium gluconate	10%. In 10 mL amps	
	10%. In 10 mL syringes	
	10%. In 10 mL single dose vials	
	10%. In 50 mL single dose vials	
	10%. In 100 mL pharmacy bulk vials	
	10%. In 200 mL pharmacy bulk vials	
Cyanokit	1 kit	
Cyanide antidote kit <i>or</i>	1 kit	
Amyl nitrite (inhalant ampules)	0.3 mL inhalant amps	
Sodium Nitrite (IV; 300mg/10ml)	30 mg/mL. In 10 mL vials	
Sodium Thiosulfate (injection solution; 100mg/ml or 250mg/ml)	10% (100 mg/mL). In 10 mL vials	
	25% (250 mg/mL). In 50 mL vials	
Deferoxamine (Desferal®; IM)	2 g vials	
	500 mg vials	
Digoxin specific antibody fragment (Digifab®, Digibind®; IV)	Digibind – 38 mg/ vial	
	Digifab – 40 mg/ vial	
Dimercaprol (BAL in oil, British anti-lewisite; IM)	10% (100mg/mL). In 3mL amps.	
DMSA (Succimer, Chemet®; capsule)	100 mg capsules	
Ethanol in D5W	5% ethanol in D5W. In 1000 mL	
	10% ethanol in D5W. In 1000 mL	

Flumazenil (Romazicon [®])	0.1 mg/mL. In 5mL vials	
	0.1 mg/mL. In 10mL vials	
Folic acid	5mg/mL. In 10mL vials.	
Folinic acid (Leucovorin [®])	3 mg/mL. In 1mL amps.	
	10mg/mL. In 5mg single-dose vials.	
	50 mg/vial	
	100mg/vial	
	350mg/vial	
Fomepizole (4-MP, Antizol [®] ; IV)	1 g/mL. In 1.5 mL vials	
Glucagon (IV)	1 mg (1 unit). In vials with 1 mL diluent	
L-Carnitine (Carnitor [®])	200 mg/mL. Single dose vial.	
Methylene blue (1% solution; IV)	1% (10 mg/mL). In 1 mL amps	
	1% (10 mg/mL). In 10 mL amps	
N-acetylcysteine (Mucomyst, [®] oral solution)	10% oral solution. In 4 mL vials	
	10% in 10 mL vials	
	10% in 30 mL vials	
	20% oral solution. In 4 mL vials	
	20% in 10 mL vials	
	20% in 30 mL vials	
N-acetylcysteine (Acetadote [®] ; IV)	20% (200 mg/mL). In 30 mL vials	
Octreotide (Sandostatin [®] ; IV)	0.05 mg/mL. In 1 mL amps	
	0.1 mg/mL. In 1 mL amps	
	0.2 mg/mL. In 5 mL vials	
	0.5 mg/mL. In 1 mL amps	
	1 mg/mL. In 5 mL vials	
Physostigmine (Antilirium [®] ; IV)	1 mg/mL. In 2 mL amps	
Phytonadione (Vitamin K ₁ , AquaMephyton [®])	1mg/0.5mL. In 0.5mL amps	
	10 mg/mL. In 1 mL amps	
Pralidoxime (2-PAM, Protopam [®] ; IV)	1g. In 20 mL vials	
Pyridoxine hydrochloride (IM/IV)	100 mg/mL. In 1 ml vials	

Name of hospital _____

Contact person, telephone number _____

Survey completed by, date completed _____

Bed size of hospital _____

Hospital trauma level _____

Please fax completed survey to the Oklahoma Poison Control Center at (405) 271-1816, by Monday, April 16, 2007.